

[https://doi.org/10.46344/JBINO.2023.v13i02\(a\).10](https://doi.org/10.46344/JBINO.2023.v13i02(a).10)

## ASSESSMENT OF THE HEALTH EDUCATIONAL PROGRAM ON THE KNOWLEDGE AND PRACTICES OF HYPERTENSIVE ADULTS IN JIZAN HOSPITAL, SAUDI ARABIA

**Ashraf Abdelrahman Elbashir., Nasreldeen Mohamed Ahmed Ali & Dr. Omer Ahmed Elrhima**

Assistant professor , Department of Nursing, Jizan University, College of Nursing Sciences, Jizan City, Kingdom of Saudi Arabia

Department of Nursing, College of Nursing Sciences, Jizan university, Saudi Arabia

Department of Basic medical science, faculty of medicine, Jizan university, Saudi Arabia

### ABSTRACT

**Background:** Hypertension is a major global public health problem. Knowledge of the risk factors and repercussions of hypertension is crucial to preventing the disease. Some populations have lower levels of knowledge. The incidence of hypertension has increased rapidly in the Saudi Arabia in the last few years. Consequently, several Saudi Arabians embraced new lifestyles and eating habits. Hypertension is the most widespread chronic public health problem globally, including Saudi Arabia.

**Objective:** The present study aimed to examine the hypertensive patient knowledge who receive health education program about hypertension.

**Methods:** This Quasi-experimental study was conducted on 150 patients with hypertension attending at clinics in Mohammed Bin Nasser hospital. Demographic and clinical data including the compliance different aspects of hypertension disease.

The data were collected from all patients and the analyzed using SPSS version 19 software, and independent *t*-test was used for significance tests at 0.05 level.

**Results:** Results of the current study, regarding correct knowledge of the studied group before and after the health education program and the different aspects of hypertension, showed that there was a significant improvement of the patients' knowledge concerning the correct knowledge of the definition of hypertension, signs and symptoms of hyper and hypotension, compliance to drug types of treatment dietary management, importance of exercise and complications of hypertension, ( $P < 0.001$ ).

**Conclusion:** The study showed that patients with hypertension in Jizan have low knowledge about hypertension. Patient education, and public enlightenment are imperative

### Keywords

Hypertension, Exercise , Knowledge, Risk.

## 1. Introduction

Saudi Arabia has approximately 35 million people (approximately half of whom are under 25 years), dispersed across 2,150,000 km<sup>2</sup> and 13 directorates (Riyadh is the capital city) [1]. It is a rapidly developing country swiftly emerging into the G20's powerful economies. Consequently, several Saudi Arabians embraced new lifestyles and eating habits[2]. Hypertension is the most widespread chronic public health problem globally, including Saudi Arabia [1]. Hypertension leads multiple complications including cardiovascular pathology, stroke, and renal failure. This controllable condition is next to smoking as a cause of mortality from avoidable cardiovascular events. Various risk factors are associated with hypertension, including growing older, being male, having fewer years of schooling, and having lower socioeconomic status. A positive correlation exists between overweight and hypertension, whereas being underweight is negatively associated with hypertension. [1].

Hypertension (HTN) is universally accepted as one of the most important risk factors in the development of cardiovascular disease (CVD), stroke and renal disease [3]. There has been a considerable increase in the prevalence of HTN in the Middle East during the last few years. In some Arab countries hypertension has become a major health problem [2]. This drastic increase in incidence of hypertension is specifically caused by a combination of man parameters, including family history [2]. Change in lifestyle, dietary habits and environmental factors [3]. Saudi Arabia is considered one of the

leading countries in Asia for the prevalence of hypertension [4]. Which may result in serious health problems in the near future if no appropriate measures are taken. The prevalence of hypertension may be a result of the marked shift in the Suadian diet, which has resulted in increased overweight and obesity [5]. Some studies have claimed the prevalence of hypertension is a cause of the tremendous increase in renal insufficiency [5]. Increased awareness, follow up and control of hypertension in industrialized countries has resulted in a decreased tendency to morbidity and mortality from cardiovascular disease [6]. To reach the level of improvement attained in developed countries, epidemiological studies on the risk factors, control methods, control levels, lifestyle, adherence to medication, and awareness will be crucial for setting control strategies in some country.

## 2. Materials and Methods

This A quasi-experimental study was conducted on patients with hypertension This study was carried out in Mohammed Bin Nasser Hospital at Jizan city. A total of 150 HNT patients were randomly selected , On January 2024. The patients participated registered hypertension patients Age 30 years and above, both sex's male and female, diagnosed with hypertension, able to communicate, visiting the clinic regularly for treatment or follow up. (Monthly or Biannually).

The questionnaire consists of 40 multiple choice questions covering different aspects of hypertension disease including

definition, types, risk factors, symptoms, complications, main aspects of self-care, and main aspects of dietary management and importance of physical activity for hypertension patients.

Revised hypertension knowledge Questionnaire was used, on a 4 levels: those who correctly answered less than 15 to 25 hypertension knowledge questions were considered to have poor knowledge, those who answered 25 to 50 of questions as having moderate knowledge, those who answered 51 to 75 of questions as having good knowledge and those who answered more than 76 as having Very good knowledge [7].

The questionnaire was filled by the researcher through using a clear Arabic language, the questionnaire includes the following sections: Patients file number, place of residence and telephone number, Sociodemographic data: including sex, age, marital status educational level, income level, Anthropometric measurements: including height, weight, body mass index, blood pressure, Hypertension related data: Duration of disease, family history of hypertension, other treatment, and

presence of complications, Health Education data: Previous health education programs and Lifestyle data: including smoking, daily and previous physical activity and dietary history.

Data was presented using descriptive statistics including frequency, percentage, mean with standard deviation (SD) and P-value of  $\leq 0.05$  was considered statistically significant for relationship investigations.

### 3. Results

Table 1. The study showed on 43 (57%) were males and 32 (42%) were females on control group is likely same above value 40 (53%) were males and 35 (427%) were females, (33%) were 40 years old. The mean  $\pm$  standard deviation of age was  $2.15 \pm 0.77$  for intervention group and  $2.01 \pm 0.78$  for control group. The majority of subjects (72%) were married in intervention group and (76%) for control group. Most of subjects (75%) were working in intervention group and (73%) on control group. There is no significant difference between intervention group and control group on the demographic characteristics at the level of (P 0.05).

**Table 4.1: The socio-demographic characteristics {Intervention group & Control groups}**

Variable	Intervention group (n= 75)		Control group (n=75)		t	p
	No	%	No	%		
<b>Age ( years)</b>						
30-	15	20	22	29	1.09	.261
40-	25	33	28	37		
50-	35	47	25	33		
Mean ± SD	2.15±0.77		2.01 ± 0.78			
<b>Sex:</b>						
Male	43	57	40	53	-0.12	.541
Female	32	42	35	47		
Mean ± SD	1.15±0.747		1.01 ± 0.58			
<b>Marital status</b>						
Single	21	28	18	24	-0.18	.40
Married	54	72	57	76		
Mean ± SD	1.95 ± 0.63		1.97 ± 0.66			
<b>Working status</b>						
Working	56	75	55	73	0.15	.653
Not working	19	25	20	27		
Mean ± SD	1.27 ± 0.45		1.25 ± 0.44			
<b>Educational level</b>						
Illiterate	15	20	17	23	-0.07	0.875
Primary	29	39	22	29		
Secondary	21	28	30	40		
University	10	13	6	8		
Mean ± SD	2.35 ± 0.95		2.33 ± 0.92			

Table 2. Shows The result of the mean value of patient's knowledge regarding definition of hypertension before attending of educational programs was  $0.33 \pm 0.43$  while the mean knowledge of the patients after the program on the same dimension was  $0.88 \pm 0.34$ , There is a significant difference ( $P = 0.002 < 0.05$ ). The mean value of patient's knowledge regarding causes of hypertension before attending of educational program was  $0.23 \pm 0.42$  while the mean knowledge of the patients after the program on the same dimension was  $0.87 \pm 0.46$ , there is a significant difference ( $P = 0.007 < 0.05$ ).

The mean value of patient's knowledge regarding signs & symptoms of hypertension before attending of

educational program was  $0.40 \pm 0.53$  while the mean knowledge of the patients after the program on the same dimension was  $0.92 \pm 0.46$ , there is a significant difference ( $P = 0.001 < 0.05$ ). The mean value of patient's knowledge regarding signs & symptoms of hypotension before attending of educational program was  $0.36 \pm 0.34$  while the mean knowledge of the patients after the program on the same dimension was  $0.93 \pm 0.48$ , there is a significant difference ( $P = 0.006 < 0.05$ ). The result of The mean value of patient's knowledge regarding Importance of compliance to types of drugs about hypertension before attending of educational program was  $0.32 \pm 0.53$  while the mean knowledge of the patients after the program on the

same dimension was  $0.65 \pm 0.45$ , there is a significant difference ( $P = 0.005 < 0.05$ ). The mean value of patient's knowledge regarding Importance of compliance to medication regimen about hypertension before attending of educational program ( $n=75$ ) was  $0.12 \pm 0.40$  while the mean knowledge of the patients after the program on the same dimension was  $0.55 \pm 0.33$ , there is a significant difference ( $P = 0.004 < 0.05$ ). The mean value of patient's knowledge regarding Importance of compliance to dietary program about hypertension before attending of

educational program was  $0.29 \pm 0.43$  while the mean knowledge of the patients after the program on the same dimension was  $0.80 \pm 0.45$ , there is a significant difference ( $P = 0.005 < 0.05$ ). The mean value of patient's knowledge regarding Importance of compliance to exercise about hypertension before attending of educational program was  $0.32 \pm 0.47$  while the mean knowledge of the patients after the program on the same dimension was  $0.87 \pm 0.46$ , there is a significant difference ( $P = 0.003 < 0.05$ ).

**Table 4.2: Distribution of correct knowledge about hypertension disease aspects between the intervention and control groups before education program (N=150)**

Item	Intervention group				t	p
	Before intervention		After intervention			
	No (%)	Mean ± SD	No (%)	Mean ± SD		
Definition of hypertension	25 (33)	$0.33 \pm 0.43$	66 (88)	$0.88 \pm 0.34$	-07.5	0.002
Causes of hypertension	17 (23)	$0.23 \pm 0.42$	65(87)	$0.87 \pm 0.46$	-12.2	0.007
Signs symptoms of hypertension	30(40)	$0.40 \pm 0.53$	69 (92)	$0.92 \pm 0.46$	-10.0	0.001
Signs symptoms of hypotension	27 (36)	$0.36 \pm 0.34$	70 (93)	$0.93 \pm 0.48$	-07.4	0.006
Importance of compliance types of drugs	24 (32)	$0.32 \pm 0.53$	49 (65)	$0.65 \pm 0.45$	-07.0	0.005
Importance of compliance to medication regimen	9 (12)	$0.12 \pm 0.40$	41 ( 55)	$0.55 \pm 0.33$	-03.2	0.004
Importance of compliance to dietary program	22 (29)	$0.29 \pm 0.43$	60 (80)	$0.80 \pm 0.45$	-07.6	0.005
Importance of compliance to exercise	24 (32)	$0.32 \pm 0.47$	65 (87)	$0.87 \pm 0.46$	-06.5	0.003

Table 3. Shows the result of the mean value of patient' correct knowledge regarding vascular complications of hypertension before attending of educational programs was  $1.81 \pm 0.39$  while the mean knowledge of the patients after the program on the same dimension was  $1.20 \pm 0.40$ , there is a significant difference ( $P < 0.05$ ). The mean value of patient's correct knowledge regarding renal complications of hypertension before attending of

educational programs was  $1.80 \pm 0.40$  while the mean knowledge of the patients after the program on the same dimension was  $1.21 \pm 0.41$ , there is a significant difference ( $P < 0.05$ ). The mean value of patient's correct knowledge regarding eye complications of hypertension before attending of educational programs was  $1.88 \pm 0.33$  while the mean knowledge of the patients after the program on the same dimension was  $1.37 \pm 0.49$ , there is a



significant difference ( $P < 0.05$ ). The mean value of patient's correct knowledge regarding cardiac complications of hypertension before attending of educational programs was  $1.85 \pm 0.36$  while

the mean knowledge of the patients after the program on the same dimension was  $1.20 \pm 0.40$ , there is a significant difference ( $P < 0.05$ ).

**Table 3. Distribution of correct knowledge about hypertension complications (n=75).**

Item	Intervention group				t	p
	Before intervention		After intervention			
	No (%)	Mean $\pm$ SD	No (%)	Mean $\pm$ SD		
Vascular complications	14 (18.7)	$1.81 \pm 0.39$	60(80)	$1.20 \pm 0.40$	7.87	0.000
Renal complications	15 (20)	$1.80 \pm 0.40$	59(78.7)	$1.21 \pm 0.41$	8.87	0.000
Eye complications	9 (12)	$1.88 \pm 0.33$	47 (62.7)	$1.37 \pm 0.49$	9.78	0.000
Cardiac complications	22 (29.3)	$1.85 \pm 0.36$	60(90)	$1.20 \pm 0.40$	8.45	0.000

Table 4. Shows the result of the mean value of patient's correct knowledge about change the diet program for hypertension disease before attending of educational programs was  $1.88 \pm 0.33$  while the mean knowledge of the patients after the program on the same dimension was  $1.33 \pm 0.47$ , there is a significant difference ( $P = 0.000 < 0.05$ ). The result of the mean value

of patient's correct knowledge about maintaining healthy dietary habits for hypertension disease before attending of educational programs was  $1.92 \pm 0.27$  while the mean knowledge of the patients after the program on the same dimension was  $1.09 \pm 0.29$ . There is a significant difference ( $P = 0.000 < 0.05$ ).

**Table 4. Distribution of correct knowledge about hypertension dietary program (n=75).**

Item	Intervention group				t	p
	Before intervention		After intervention			
	No (%)	Mean $\pm$ SD	No (%)	Mean $\pm$ SD		
Change in diet program	11 (14.6)	$1.88 \pm 0.33$	50(66.7)	$1.33 \pm 0.47$	7.58	0.000
Maintenance healthy dietary habits	6 (8.0)	$1.92 \pm 0.27$	68(90.7)	$1.09 \pm 0.29$	16.9	0.000

Table 5. Show The most common source of knowledge related to information about hypertension for the patients in (Intervention group) was found to be Friends/Relatives (41.3%) followed by Mass media: Television/Newspaper 30.7%). In

only 21.3% cases the knowledge was obtained from medical and paramedical professionals. And (6.7) from others sources, the percentage is similar for the control group. There is no significant difference ( $p > 0.05$ ).

**Table 5. Distribution of Sources of information knowledge regard hypertension disease.**

Source of Knowledge	Intervention group (n=75)		Control group (n=75)	
	No	%	No	%
Friends & relatives	31	41.3	29	38.7
Media (TV, Newspaper)	23	30.7	28	37.3
Medical professionals ( physician, nurses, nutritionist)	16	21.3	11	14.7
Others	5	6.7	7	9.3

#### 4. Discussion

Regarding concerning correct knowledge of the intervention group in table 2 about definition of hypertension signs & symptoms, causes and etc....., results of the study showed more half of the hypertensive patients 68.0% had average or poor knowledge about hypertension, whereas only 16.0% patients had very good knowledge about the hypertension on the same dimension. It was improve after educational intervention to (73.3). This finding is supported by Williams MV et al [8]. In their study in 402 hypertensive patients, also found 189 patients (47.02%) did not have adequate knowledge about hypertension. Similar inadequacy of knowledge, awareness and practice of hypertension among Indian patients has been reported by (Hemant Mahajan) [9]. Importance of compliance to (drugs, medication regimen, dietary program and exercise) it was improve after educational intervention. However, the source of knowledge was significantly increased after the educational intervention, demonstrating the beneficial effects of education on the patient's knowledge. This result supported by (Falaschetti E, et al) He said hypertension control signifies a greater need to increase the awareness of hypertension related information among

the patients [10]. The possible reasons to lower knowledge may be because of lower literacy, inappropriate perception of medical advice, irregular sources of health related information, or inadequate counselling regarding hypertension possibly due to skewed doctor patient ratio in government run hospitals [11]. Regarding sources of knowledge for hypertension the patients reported to have derived their knowledge about Hypertension majorly from non-medical sources like friends/relatives and mass media communications. (72%) and only (21.3%) of sources were from medical professionals like doctors, specialist paramedical staff, which form the more reliable source to provide health related information. Therefore the patient has to be alerted to scrutinize the information received, from their doctors and work in collaboration with health providers to get valid information. The knowledge from such unreliable sources may be the cause of lower awareness among the population. Considering the influence of mass media on the population, a possible alternative to increase awareness may be by means of delivery of such information using mass media. This findings supported by (Hroscikoki MC, et al) [11]. Reported that positive role of pharmacist mediated

counselling of hypertensive patients, regarding risk factor and associated comorbidities, while some other studies suggest that knowledge transferred from medical staff is an important factor in inducing patient to comply with lifestyle modification [11]. Nevertheless; low counselling rates were reported in similar studies [12, 13]. About knowledge about lifestyle modifications (change the diet program and maintaining continuous follow up) study showed most of hypertensive patients knowledge about lifestyle modifications (32.0%) before education program, It was improve after educational intervention to (74.7). These result supported by many authors (Chiu CW, Ostovan MA) They said Patient's involvement in self-monitoring and management, together with continuous follow up has also been recommended by others [14, 15]. Similarly, Wang YR et al. emphasized that the most important points for BP control were lifestyle modifications, home BP monitoring, reinforcement of healthy behaviours, and continuous follow up [16]. In (Aubert et al) study, most patients believed that salty diet, obesity and smoking are important factors in hypertension. They mentioned that physical activity and exercise are very important factor in hypertension management [17]. Regarding hypertension complications the study showed patient's knowledge (15%) before attending of educational programs it was improve after educational intervention to (75%). These findings were supported by previous study results shown a positive relationship between patients' knowledge about the hypertensive complications and

adherence [18]. In addition, they are aware of hypertension complications and advantages of lowering blood pressure. Most cases agreed that decreasing blood pressure (even a little bit) could be effective on health and decrease there complications [19].

The strengths of this study include the study conducted in developing countries with limited resources. The study limitations were; the study was conducted as a descriptive study; interventional studies will yield more useful results if conducted on more sample with complete randomization all over the country.

## 5. Conclusion

After the analysis of the study variables the researcher justify the findings and conclude that: The results from this short educational intervention program on hypertensive patients in Mohammed Bin Nasser Hospital at Jizan City indicated that knowledge changes in a positive results in all the variables related to patients knowledge. In the current study, the majority of the patients did not have sufficient knowledge to complications of hypertension at the pre-test phase, education program effect and improve their knowledge about complication post-test there is a significant difference ( $P < 0.05$ ). Also in the current study, the majority of the patients get information about hypertension and its management from non-medical sources (from friends/Relatives (41.3%) followed by mass media: Television/Newspaper (30.7%). In only 21.3% cases the knowledge was obtained from medical and paramedical professionals. In the current study, the most of the patients did not have controlled



blood pressure at the pre-test phase only 32% of patients controlled blood pressure, Post-test after intervention 92% of patients controlled blood pressure and good knowledge to adjust their blood pressure lifestyle changes involving dietary and exercise being effective in significant decrease in weight, and effective in improving patient's knowledge.

### Recommendations

After obtaining the study findings based on the conclusion, the researcher recommended that: The educational intervention program should be developed in hypertension clinics in Jizan hospital in Saudi Arabia. Since hospitals & primary health care centers providers have a better chance to meet with patients, promotion of hypertension educational intervention programs by these centers will increase the effectiveness of hypertensive therapy and will delay the onset or the progression of complications, improve the quality of life for hypertensive patients and reduce the associated medical costs. Educating the patient's family about hypertension disease meal planning and their dietary management and the relationship between obesity and the chance of disease occurrence. Educating the family about the importance of involvement of hypertensive patient s food with family food. The ideal treatment would consist of a comprehensive and multidisciplinary Hypertension disease team. (Physician Nurse, Nutritionist, Pharmacist, Social worker, foot specialist and others). Training of the hypertensive care team on the management of hypertension and how to educate the hypertensive patients Establishment of

regional records for hypertensive patients in order to facilitate health care and health education for them.

### References

1. Herzallah HK, Antonisamy BR, Shafee MH, Al-Otaibi ST. Temporal trends in the incidence and demographics of cancers, communicable diseases, and non-communicable diseases in Saudi Arabia over the last decade. *Saudi Med J*. 2019;40(3):277.
2. Herzallah HK, Antonisamy BR, Shafee MH, Al-Otaibi ST. Temporal trends in the incidence and demographics of cancers, communicable diseases, and non-communicable diseases in Saudi Arabia over the last decade. *Saudi Med J*. 2019;40(3):277–286.
3. Gosadi IM. Assessment of the environmental and genetic factors influencing prevalence of metabolic syndrome in Saudi Arabia. *Saudi Med J*. 2016;37(1):594.
4. Saudi Cancer Registry. Cancer Incidence Report, Saudi Arabia 2015. Available from: <https://nhic.gov.sa/eServices/Documents/E%20SCR%20final%206%20NOV.pdf>. Accessed 8 May, 2020.
5. Gosadi IM. National screening programs in Saudi Arabia: overview, outcomes, and effectiveness. *J Infect Public Health*. 2019;12(5):608–614.
6. Tyrovolas S, El Bcheraoui C, Alghnam SA, et al. The burden of disease in Saudi Arabia 1990–2017: results from the Global Burden of Disease Study 2017. *Lancet Planetary Health*. 2020;4(5):e195–e208.
7. Saudi Ministry of Health. World Health Survey; 2019. Available from: <https://www.moh.gov.sa/en/Ministry/Statistics/Population-Health-Indicators/Documents/World-Health-Survey-Saudi-Arabia.pdf>. Accessed May 6, 2022.
8. Williams MV, Baker DW. Relationship of functional health literacy to patient's

- knowledge of their chronic disease. A study of patients with hypertension and diabetes. *Arch Intern Med*, 1998 Jan 26; 158 (2): 166-72.
9. Mahajan H, Kazi Y, Sharma B, Velhal GD. Assessment of KAP, Risk Factors and Associated Co-Morbidities in Hypertensive Patients, (Sep-Oct. 2012), PP 06-14.
  10. Falaschetti E, Chaudhury M, et al. Continued improvement in hypertension management in England: results from the Health Survey for England 2006. *Hypertension*. 2009 Mar; 53 (3): 480-6.
  11. Hroschicki MC, Solberg LI, et al: Challenges of change a qualitative study of chronic care model implementation. *Ann. Fam. Med*. 2006 Jul-Aug; 4 (4): 317-26.
  12. Prochaska J. DiClemente C. *The Trans theoretical Approach: Crossing traditional boundaries of therapy*. Homewood Il: Dow Jones-Irwin, (1984).
  13. Stange KC, Flocke SA, et al. Direct observation of rates of preventive service delivery in community family practice Preventive. 2000 Aug; 31 (2 Pt 1): 167-76.
  14. Chiu CW, Wong FK. Effects of 8 weeks sustained follow-up after a nurse consultation on hypertension: a randomized trial. *International journal of nursing studies*. [Randomized Controlled Trial Research Support, 2010.
  15. Ostovan MA, Zibaeenezhad MJ, et al The impact of education on weight loss in overweight and obese adults. *International cardiovascular research journal*. 2013 Sep; 7 (3): 79-82.
  16. Wang YR, Alexander GC, Stafford RS. Outpatient hypertension treatment, treatment intensification, and control in Western Europe and the United States. 2007 Jan; 167 (2): 141-7.
  17. Aubert L, Bovet P, Gervasoni JP, Rwebogora A, Waeber B, et al. Knowledge, attitudes, and practices on hypertension in a country in epidemiological transition. 1998 May; 31 (5): 1136-45.
  18. Ghembaza MA, Senoussaoui Y, Tani MK, Meguenni K. Impact of patient knowledge of hypertension complications on adherence to antihypertensive therapy. 2014; 10 (1): 41-8.
  19. Egan BM, Lackl and DT, Cutler NE. Awareness, knowledge, and attitudes of older Americans about high blood pressure: implications for health care policy, education, and research. 2003 Mar 24; 163 (6): 681-7.