

## IMPACT OF URINE NITRATE ON ATTACHED EAR LOBE

Muhammad Imran Qadir, Fakhra Batool, Asma Rasheed\*

Institute of Molecular Biology and Biotechnology, Bahauddin Zakariya University, Multan,

### ABSTRACT

Objective of present study was to co-relate the nitrate in urine with joined ear lobe. The most likely cause of nitrites in urine is UTIs. A total 90 subjects participated in this project. We took dipstick for the test of urine nitrate. We determined the presence of nitrate in the urine by taking the sample and dipped the test stick into the sample and noticed the change of color which showed that the nitrate is present. If no change occurs then nitrate will not be present. It was concluded from the present study that the males and females who have attached ear lobe have no urine nitrate but the males and females with free ear lobe have urine nitrate

**KEY WORDS:** *Urine nitrate, joined ear lobe, test strips*



## INTRODUCTION

When nitrites in urine occur then it because the urinary tract infection and this is due the infection caused by bacteria. Nitrites are derivative of waste of nitrogen and this waste made the food of bacteria and breaks it into nitrates then it appears in the urine. When the nitrates appear in the urine then antibiotics are used. A doctor will acquire the individual's medical record, any hypersensitivity, and the kind of microorganisms, if it is identified, when recommends an antibiotic. The majority expected reason of nitrites in urine is UTIs. Particularly persons in the bladder and urethra are simple to indulgence and will apparent up within a week of captivating antibiotics.

There are two types of ear lobe which are attached ear lobe and free ear lobe and it can be controlled by the gene. Further appearance of vital inheritance unattached ear lobes is used. The representation is that earlobes can be alienated into 2 types which are joined and free ear lobe and that thorough innate substance manage the accomplishment. The dominant is free ear lobe. On the other hand, numerous of the papers on earlobe inheritance have demonstrate that there are many people with middle earlobes El Kollali divided the

earlobes into 3 types which were based on the point of view which were liable, exact, or not susceptible.

## MATERIALS AND METHOD


A total 90 subjects participated in this project. We took dipstick for the test of urine nitrate. We determined the presence of nitrate in the urine by taking the sample and dipped the test stick into the sample and noticed the change of color which showed that the nitrate is present. If no change occurs then nitrate will not be present.

## Statistical analysis

The statistical analysis was done by the m stat.

## RESULTS AND DISCUSSION

This graph shows that 54.5% males have attached ear lobe but none of them contain urine nitrate. The 45.4% males have free ear lobe but 9.69% have urine nitrate and 36.3% have no urine nitrate. The 29.8% females have attached ear lobe and none of them contain urine nitrates. The 70.1% females have free ear lobe and 1.49% have urine nitrate and 68.6% females with free ear lobe have no urine nitrate.



## CONCLUSION

It was concluded from the present study that the males and females who have attached ear lobe have no urine nitrate but the males and females with free ear lobe have urine nitrate.

## REFERENCE

**Qadir MI, Malik SA** (2010) Comparison of alterations in red blood cell count and alterations in hemoglobin concentration in patients suffering from rectal carcinoma undergoing 5-fluorouracil and folic acid therapy. *Pharmacologyonline*, 11 3: 240-243.

**Qadir MI, Noor A** (2018) *Anemias. Rare & Uncommon Diseases*. Cambridge Scholars Publishing. Newcastle, England. ISBN: 978-1-5275-1807-0.

**Qadir MI, Javid A** (2018) Awareness about Crohn's Disease in biotechnology students.

*Glo Adv Res J Med Medical Sci*, 7(3): 062-064.

**Qadir MI, Saleem A** (2018) Awareness about ischemic heart disease in university biotechnology students. *Glo Adv Res J Med Medical Sci*, 7(3): 059-061.

**Qadir MI, Ishfaq S** (2018) Awareness about hypertension in biology students. *Int J Mod Pharma Res*, 7(2): 08-10.

**Qadir MI, Mehwish** (2018) Awareness about psoriasis disease. *Int J Mod Pharma Res*, 7(2): 17-18.

**Qadir MI, Shahzad R** (2018) Awareness about obesity in postgraduate students of biotechnology. *Int J Mod Pharma Res*, 7(2): 14-16.

**Qadir MI, Rizvi M** (2018) Awareness about thalassemia in post graduate students. *MOJ Lymphology & Phlebology*, 2(1): 14-16.

**Qadir MI, Ghalia BA** (2018) Awareness survey about colorectal cancer in students of M. Phil Biotechnology at Bahauddin Zakariya University, Multan, Pakistan. Nov

Appro in Can Study, 1(3): NACS.000514.2018.

**Qadir MI, Saba G** (2018) Awareness about intestinal cancer in university student. Nov Appro in Can Study, 1(3): NACS.000515.2018.

